

Evaluation of Folic Acid level in Serum of Smokers and Non Smokers Patients with Periodontal Disease: A Case-Control Study

**Ghasaq Asim Abdul-Wahab¹, Alyamama Mahmood Alwan², Sarah Ihsan Al-Karawi³,
Tameem Jassim Khuder⁴**

¹Oral Surgery and Periodontology Department / College of Dentistry /Mustansiriyah University,
E-mail: Ghasaq.a.abdulwahhab@uomustansiriyah.edu.iq, ORCID ID: <http://orcid.org/0000-0001-6654-9416>.

²Oral Surgery and Periodontology Department/ College of Dentistry /Mustansiriyah University,
E-mail: alyamamahmahmood@uomustansiriyah.edu.iq, ORCID ID: <https://orcid.org/0000-0002-4601-7516>.

³Oral Surgery and Periodontology Department / College of Dentistry /Mustansiriyah University,
E-mail: saraihsan@uomustansiriyah.edu.iq, ORCID ID: <https://orcid.org/0000-0002-7994-9635>.

⁴Prosthodontics Department / College of Dentistry /Mustansiriyah University,
E-mail: tameem.kh@uomustansiriyah.edu.iq, ORCID: <https://orcid.org/0000-0002-7630-599X>.

Received 03/11/2024

Accepted in revised form 27/12/2024

Published 30/12/2024

Abstract

Objectives: Periodontitis is a complex disease, the reason of which remained unclear. Folic acid belongs to B complex vitamins. It is vital for the formation and division of new cells, as it serves as a crucial cofactor in DNA synthesis. Without the presence of folic acid, epithelial cells fail to divide correctly. Folic acid levels can be influenced by a number of factors, one of which is cigarette smoking. This study aimed to measure serum concentration of folic acid among patients with periodontitis associated to their smoking behaviors. **Materials and Methods:** The study included 75 subjects; divided in the following groups: group-I: 25 Smokers with periodontitis, group-II: 25 Non-smokers with periodontitis and group-III: 25 Non-smoker with healthy periodontium (control group). Clinical parameters were recorded including: Plaque Index(PLI), Gingival Index GI, Bleeding On Probing (BOP), Probing Pocket Depth(PPD) and Clinical Attachment Loss(CAL). Blood samples were taken and evaluated in the laboratory for folic acid levels by enzyme-linked immunosorbent assay ELISA technique. **Results:** Significantly low concentration of folic acid encountered in serum of smokers with periodontitis as compared to healthy non-smoker controls ($P < 0.001$). Additionally, smokers with periodontitis showed significantly higher PLI, PPD and CAL means than non-smokers with periodontitis

Conclusion: The results of the present study indicate that folic acid levels were decreased significantly in smokers with periodontitis. Smoking has a negative impact on folic acid level and periodontal health.

Key Words: Periodontitis, Serum, Folic acid, Cigarette Smoking, ELISA.

Introduction

Periodontitis is a common inflammatory disease affecting the tooth-supporting tissues, started by specific pathogens resulting in persistent deterioration of the alveolar bone and periodontal ligament. Periodontal tissue destruction happens by an exaggerated host immune-inflammatory response.¹ Around 40% and 90% of individuals worldwide affected by periodontitis, resulting in the most common epidemics in the world.²

Periodontal health depends on a precise equilibrium among the host, bacteria, environment, and different microorganisms. Periodontal disease results from complex interactions between the host factors and infectious agents. Many risk factors such as environmental, acquired, and genetic could change disease expression and may impact its onset, progression, and treatment response.³

Tobacco smoking is a widespread habit that has been established as linked to cancer. Also, it may have a role as an environmental factor linked to a higher prevalence and severity of periodontal disease, while also adversely affecting nutritional levels. Nutrition is a modified factor that influences the host's immune response as well as integrity of the mouth cavity's soft and hard tissues.^{4,5}

The presence of essential nutrients in the host significantly influences periodontal health. Among the human body's most active tissues are the epithelium of the dentogingival junction and the adjoining connective tissues. Periodontium health requires sufficient proteins, mineral salts, lipids, carbs, and vitamins. A chronic deficit in any of these nutrients may cause periodontal tissue pathology.^{6,7}

Folic acid (named as vitamin B9 or folacin), along with folate (the naturally occurring type) are type of vitamin B9, which is water-soluble. It's crucial for biosynthesis of nucleotide and the homocysteine remethylation. Folic acid plays an important role during phases of rapid cellular division and development. Folate and folic acid terms originated from the Latin-word; folium; which means "leaf".³

Cigarette smoking is among those factors that negatively impact folic acid levels.⁸ Cigarette smoke contains nitric oxide, superoxide and various species of reactive oxygen. Smoking's negative consequences may arise from oxidative damage to endothelial cells.⁹ In inflammatory reactions; the production of significant amounts of nitric oxide and superoxide, their interaction results in reactive nitrogen species generation, including the peroxynitrite anion, known as a toxic

byproduct of nitric oxide together with superoxide. Peroxynitrite-induced tissue injury due to exaggerated local immune response and inflammatory cells migration. The bioactive form of folic acid (5-Methyl Tetrahydro-folic acid) has a demonstrated ability to decrease superoxide generation, thereby manifesting an antioxidant capacity. Conversely, an oxidant and a free radical, which induce damage to DNA and membrane, found in elevated levels in tobacco smoke and could potentially contribute to the reduction of folic acid levels. Cigarette smoke comprises organic the nitrite nitrous oxide (NO), cyanates, and isocyanates which react with co-enzymes of B12 vitamins and folic acid to transform them into molecules that are not biologically active. In addition to the negative impacts of smoking on immunity and interaction between host and bacteria, it may contribute to its damaging effect on periodontal health.⁹ Moreover, smoking adversely influences folic acid mechanisms of action. The insufficient amount of this vitamin could hinder the formation and evolution of the periodontal tissues, as well as the beginning and advancement of periodontal diseases.³ Thus; the purpose of this study was to assess and juxtapose serum the amount of folic in smokers and non-smokers, with and without periodontitis, and to investigate result of folic acid levels on periodontal health status.

Subjects, Material and Methods:

Study design: Case-control study.

Sample Size: The sample size was calculated with G*Power 3.1.9.7 at a power of 0.9 and margin of error of 5% (0.05).¹⁰ The calculated sample size for periodontitis

group is 50, while it will be 25 for healthy controls. So, the total sample size would be 75 (25 control, 50 smokers and non-smokers with periodontitis).

Subject Population:

The study included seventy-five individuals with age range of 30- 55 years old; consisting of fifty participants with generalized stage II periodontitis divided into two groups each group consist of twenty-five patients with stage II periodontitis smokers and non-smokers; and twenty-five periodontally healthy non-smokers volunteers. The subjects involved were possessed at least 20 natural teeth, showed the desire with the ability for signed a consent form and systemically healthy without any diseases or conditions that affect the periodontium. Smokers' patients, who smoked for five years or more, between 5 and 10 cigarettes\day.

The criteria of generalized stage II periodontitis were defined as the presence of more than 30% of sites with pocket depths less than 5 mm and/or 30% of sites with clinical attachment loss > 4 mm. In healthy individuals, less than 30% of sites had a gingival index less than 1, and no CAL.¹¹ The participants with systemic diseases that might affect the development or treatment of periodontal disease, those who had received antibiotics in the last three months, and individuals who used any vitamins or nutrition supplements in the past six months were excluded. The ethical committee of the College of Dentistry approved this study under the reference number of REC143 and study number MUOSU-202112. Every participant signed a written informed

consent form before being enrolled in the study.

Clinical Parameters

Clinical parameters PLI, GI, PPD and CAL were measured for all the participants. The recordings were done at 6 sites per tooth (mesio-buccal, mid-buccal, disto-buccal, disto-lingual, mid-lingual and mesio-lingual) for all teeth except third molars. The records were taken using William probe.

Serum Analysis

Samples of blood were obtained from fasting participants, as recent food consumption can significantly elevate serum folate levels.¹² Under aseptic conditions; 5 milliliters of intravenous blood were obtained from all participants; the blood sample was then placed to plain tube to clot for 1 hour, and then at room temperature the sample centrifuged at 3300 rpm for 5 minutes. The clear serum sample was then separated and stored at -20°C until analysis. Then the samples were sent to the laboratory to assess serum levels of folic acid, utilizing ELISA technique.

Data Collection and Statistical Analysis

The mean and standard deviation (SD) of plaque Index, gingival index, PPD and CAL were calculated. Also, serum levels of folic acid's mean and standard deviation were calculated. All study groups follow a normal distribution therefore a parametric test of one-way ANOVA; Least significant difference test (LSD) and T-test was conducted for intergroup comparisons. Data was collected and analyzed via IBM®

SPSS® version 27 with a P-value of <0.05 considered as significant.

Results:

According to the demographic results, the mean age of Group I was 41.72 ± 7.34 years, Group II was 39.76 ± 7.06 years and Group III was 38 ± 7.29 years respectively. The clinical periodontal parameters (PI, GI, PPD and CAL) were higher in group I followed by group II and III respectively.

Folic acid values demonstrated least levels in smokers with periodontitis (group I) as compared to non-smokers with periodontitis (group II) 4.004 ng/ml, 9.572 ng/ml respectively; while the controls (group III) showed highest level of folic acid (16.8 ng/ml) (Table I).

For intergroup comparisons there were highly significant differences ($P < 0.0001$) both between and within study groups for serum folic acid level, PLI, GI, PPD and CAL (Table II).

Upon intergroup comparison of three study groups, significant differences encountered in mean of serum folic acid between group I and II; group I and III; group II and III ($p < 0.0001$) suggesting its level being lower in smokers. Likewise, highly significant differences in mean of PLI and GI was found between (group I and group III) and between (group II and group III) ($P < 0.0001$). While and nonsignificant differences between group I and II regarding PLI and GI ($P > 0.05$) as shown in (Table-III).

According to Pearson's rank correlation (r) of clinical parameters and levels of Folic

Acid in study groups (Table- IV); the result reflect an inverse significant association were demonstrated between serum folic acid

and PLI in group II; and between folic acid and GI, PPD, CAL in group I and II (P<0.05).

Table-I: Mean and Standard Deviation of Age, Periodontal Parameters, and Serum Folic Acid of the Study Groups

Parameters	Group I	Group II	Group III
	Mean \pm SD	Mean \pm SD	Mean \pm SD
Age	41.72 \pm 7.346	39.76 \pm 7.061	38 \pm 7.292
PLI	2.376 \pm 0.32	2.13 \pm 0.41	0.30 \pm 0.16
GI	2.12 \pm 0.67	2.56 \pm 0.287	0.37 \pm 0.16
PPD	5.27 \pm 0.65	4.28 \pm 0.38	-
CAL	7.28 \pm 0.85	4.45 \pm 0.57	-
FolicAcid (ng/ml)	4.004 \pm 1.14	9.572 \pm 1.48	16.8 \pm 1.08

SD: Standard Deviation; PLI: Plaque Index; GI: Gingival Index; PPD: Probing Pocket Depth; CAL: Clinical Attachment Loss; ng/ml: nanogram/milliliters.

Table-II: ANOVA Test for Serum Folic Acid and Clinical Periodontal Parameters Among Study Groups

Index	ANOVA Test	SS	df	MS	F- test	P-value	Significance
Folic Acid	Between Groups	2058.20	2	1029.10	662.74	<0.00001	*HS
	Within Groups	111.8	72	1.55			
	Total	2170.00	74				
PLI	Between Groups	64.24	2	32.12	323.26	<0.0001	*HS
	Within Groups	7.15	72	0.09			
	Total	71.40	74				
GI	Between Groups	66.84	2	33.42	175.645	<0.0001	*HS
	Within Groups	13.70	72	0.19			
	Total	80.548	74				
PPD	T- test	6.4791			<0.0001		*HS
CAL	T- test	13.6765			<0.0001		*HS

HS: highly Significant; Sum of Squares: SS; DF:Degree of Freedom;MS: Mean Square and statistical expressions: F and P; PLI: Plaque Index; GI: Gingival Index; PPD: Probing Pocket Depth; CAL: Clinical Attachment Loss.

Table-III: Least Significant Difference Test (LSD) to Compare the Means of Serum Folic Acid Level and Clinical Periodontal Parameters Among Study Groups

Parameters	Groups	MD	SE	P- value	Significance
Folic Acid	GI- GII	5.57	22.34	0.000	HS
	GI- GIII	12.80	51.34	0.000	HS
	GII- GIII	7.23	29.00	0.000	HS
PLI	GI- GII	0.25	3.90	0.0198	S
	GI- G III	2.07	32.91	0.000	*HS
	G II- GIII	1.83	29.01	0.000	*HS
GI	GI- GII	0.44	5.00	0.0020	S
	GI- GIII	1.75	20.04	0.000	*HS
	GII- GIII	2.18	25.04	0.000	*HS

S: Significant, NS: Non-significant, * HS: Highly Significant ; PLI: Plaque Index; GI: Gingival Index; PPD: Probing Pocket Depth; CAL: Clinical Attachment Loss; MD :Mean Difference, SE: Standard Error, P: P-value.

Table-IV: Spearman's Rank Correlation (r) of Serum Folic Acid and Clinical Periodontal Parameters Among Study Groups

Groups and Parameters		Group I	Group II	Group III
		Folic Acid		
PLI	r	0.209	-0.389	-0.068
	P	0.316	0.054*	0.7467
GI	r	-0.938	-0.609	-0.058
	P	<0.0001**	0.0012**	0.7830
PPD	r	-0.671	-0.382	-
	P	0.0002 *	0.053*	-
CAL	r	-0.821	-0.50	-
	P	<0.0001**	0.0109*	-

r: Spearman's Rank Correlatio; * Significant; ** Highly Significant; ; PLI: Plaque Index; GI: Gingival Index; PPD: Probing Pocket Depth; CAL: Clinical Attachment Loss.

Discussion:

Periodontal disease is a multifactorial condition characterized by complex etiology and wide phenotypic variation. Alongside the primary factor of bacterial plaque, inherited and social conditions, as well with the individual's behaviors, play crucial supporting roles. The increasing prevalence of smoking addiction and its related mortality as well as mortality renders

smoking a serious public health risk.¹³ However, the exact mechanism through which smoking affects is still unknown. It is unclear if smoking induces systemic effects on the periodontium or local effects that promote periodontal disease.^{14,15}

The age group of 30-55 years was selected for the present study because patients in this age range are most likely to indulge in

deleterious habits like cigarette smoking. According to a study done by Singh *et al* (2014), The prevalence of smoking was greater among older persons, individuals residing in rural areas, the uneducated, and the impoverished, in contrast to the younger population and those living in metropolitan areas¹⁶. Also, the majority of epidemiological research found that both the severity and prevalence of chronic periodontitis are higher at this age; female patients were excluded from this study as recruiting women who acknowledge that they smoke would ultimately be challenging.

The periodontal parameters of the study groups were recorded to assess the oral hygiene status. On intergroup comparison, there was a statistically significant and highly significant difference between all groups ($p < 0.001$). The results of the present study indicated that plaque accumulation was minimal in the healthy control group (0.30 ± 0.16), followed by non-smokers with periodontitis group (2.13 ± 0.41), then smokers with periodontitis group (2.376 ± 0.32). The oral hygiene status indicated by plaque levels correlates with prior studies by Torrungruang *et al* in 2005; who shown significantly higher plaque levels in smokers¹⁷. While contradicting to findings of other previous studies.^{14,18,19}

The difference between the mean gingival index in all study groups was statistically highly significant ($p < 0.001$). On intergroup comparison, a statistical significant difference was observed between Group I and Group II ($P = 0.002$). The results of this study indicate that the GI was greater in the non-smokers' group in contrast to both the smokers and the healthy controls groups.

These results agree with previous studies by Jalayer *et al* (2015)²⁰ who have shown lower levels of the gingival index. This could be due to tobacco products, which impact vascular inflammation. Contrary to these results, the findings of researchs performed by Johanssen *et al* in 2005²¹, and Arowojolu *et al* in 2013.²²

Highly significant differences were also observed in the mean of PPD, and CAL scores between smokers and non-smokers with periodontitis. The mean of PPD in (smokers with periodontitis group) was 5.27 ± 0.65 and in (non smokers with periodontitis group) was 4.28 ± 0.38 . The difference between the mean of PPD was statistically highly significant ($p < 0.001$) as the smokers exhibit a higher scores of PPD compared to non-smokers group. The results of our study agree with a study done by Rahul *et al*, 2024; Ragghianti *et al.*, 2004; and Velidandla *et al*; 2019.^{14,23,24}

Concerning CAL, the difference between the mean of CAL in smokers and non-smokers was statistically highly significant ($p < 0.001$). The smokers group with periodontitis record a higher mean of CAL (7.28 ± 0.85); while the non-smokers recorded (4.45 ± 0.57). This highlights that the clinical attachment level was more in the smokers as compared to the non-smoker's group. CAL levels are a consequence of an inflammatory burden from the past into the present, while probing pocket depth levels reflect the ongoing pathophysiological status of periodontitis. Our study's findings may be related to modifications in subgingival plaque structure, and subgingival bacteria virulence and host response, which increase periodontium deterioration and bone

resorption. Tobacco's nicotine can harm collagen tissues by increasing the production of collagenase, inhibiting the growth of gingival fibroblasts and decreasing the synthesis of collagen and fibronectin which ultimately compromises periodontal health. The results of our study agreed with Rahul et al, 2024; Torrungruang, 2005, Calsina et al, 2002, Susin et al, 2004; Rudziński and Banach, 2011 studies.^{14,17,19, 25, 26}

Recent research suggests that nutritional habits and dental health have potential link in older persons.^{27,28,29} Nutrition greatly impacts the health of the periodontium, and its deficiency can alter the presentation of the main etiological factors. Periodontium, one of the body's most dynamic tissues, with its sustainability reliant on a sufficient nutrients supply.³⁰

As folic acid, considered as a key vitamin within the B vitamin group, is recognized as a hemo-cytopoietic vitamin and plays a role in growth.³¹ Folic acid deficiency represents the most prevalent nutrient deficiency globally and it correlates with elevated oxidative stress, malfunctioning endothelial cells, genomic disorder, weakened DNA repair, and apoptosis. It is has been demonstrated to be associated with various diseases of humans, including periodontal diseases. However, very little has been discovered about how smoking affects folic acid level in chronic periodontitis patients. Clinical studies suggest that folic acid supplementation reduces gingival inflammation, characterized by redness, bleeding, tenderness and exudates.³²

According to this study the folic acid mean in smokers with periodontitis was 4.004 ± 1.14 ng/ml, while non-smokers were 9.572 ± 1.48 ng/ml and controls were 16.8 ± 1.08 ng/ml. Intergroup comparisons indicated highly significant differences; concentrations of serum folic acid were notably less in smokers with chronic periodontitis group versus nonsmokers with similar periodontal destruction group and controls with healthy periodontium group. This aligns with the research conducted by Erdemir and Bergstrom^{8,33}, Sumona et al³⁴ Yu et al.³⁵

Low folic acid levels may result from the reaction between isocyanates and cyanates together with folic acid, turning it inactive biologically and consequently reducing its levels. However, it may also result from insufficient intake of nutrients, poor absorption or metabolic dysfunction.

The correlation between the clinical parameters and folic acid was a strong negative correlation was found with PLI in non-smokers group and negative significant correlation with GI, PPD and CAL in both smokers and non-smokers with periodontitis, these inverse correlations signifying more folic acid levels were associated with reduced loss of attachment in patients with chronic periodontitis, regardless of smoking status. It can be inferred that increased folic acid levels may lead to decreased destruction of periodontium. Consequently, cigarette smoking habit, a known risk factor for periodontal diseases, could magnify the destruction resulting from folic acid deficiency. These results were in accordance with findings of Sumona et al, 2011.³³

While smoking cessation is the preferred goal, it is not always achievable; thus, any strategy aimed at minimizing the harmful effects of smoking is valuable. Smokers with inadequate folic acid levels may benefit from enhanced dietary intake or supplementation of folic acid. A longitudinal study might explain the exact period of the deficient state and disease initiate, and providing stronger proof. Thus, longitudinal and interventionist studies in more epidemiological population will be required to determine the link between periodontal disease, folic acid, and smoking.

Conclusion:

The current research suggests that one significant environmental risk factor associated with further periodontal destruction is cigarette smoking. The escalated progression and excessive deterioration of periodontal support in later life largely influenced by excessive smoking in youth. Also, smoking can have negatively impacted hematological parameters like serum folic acid levels.

These findings highlight the necessity for preventive strategies targeting both younger and older individuals increasing folic acid intake by diet or folic acid supplementation may beneficial to dental health.

Acknowledgements

The authors would like to thank Mustansiriyah University (www.uomustansiriyah.edu.iq), Baghdad, Iraq, for their support in the present work.

Conflict of interest

The authors reported that they have no conflicts of interest.

Ethical approval: The study was approved by the institutional ethical committee.

References:

1. Hans M, Malik PK, Hans VM, Chug A, Kumar M. Serum levels of various vitamins in periodontal health and disease- a cross-sectional study. *J Oral Biol Craniofac Res.* 2023 Jul-Aug;13(4):471-475. doi: 10.1016/j.jobcr.2023.05.009.
2. Abdul Wahab G.A., Alwan A.M., Al Karawi S.I. 2024. Evaluation of Serum Interleukin 33 Level in Iraqi Patients with and without Periodontal Disease, Clinical, Cosmetic and Investigational Dentistry, 201-207, DOI: 10.2147/CCIDE.S464951.
3. George JP, Shobha R, Lazarus FJ. Folic acid: A positive influence on periodontal tissues during health and disease. *Int J Health Allied Sci* 2013;2:145-52
4. Nunn ME. Understanding the etiology of periodontitis: An overview of periodontal risk factors. *Periodontol 2000* 2003;32:11-23.
5. Seymour RA, Heasman PA. *Drugs, diseases and the periodontium.* New York; Oxford University Press; 1992.
6. Apatzidou DA, Riggio MP, Kinane DF. Impact of smoking on the clinical, microbiological and immunological parameters of adult patients with periodontitis. *J Clin Periodontol* 2005;32:973-83.
7. George, Joann & Shobha, R & Lazarus, Flemingson. (2013). Folic acid: A positive influence on periodontal tissues during health and disease. *International Journal of*

- Health & Allied Sciences. 2. 10.4103/2278-344X.120582.
8. Erdemir EO, Bergstrom J. Relationship between smoking and folic acid, vitamin B12 and some hematological variables in patients with chronic periodontal disease. *J Clin Periodontol* 2006;33:878-84.
 9. Enwonwu CO, Sanders C. Nutrition: Impact on oral and systemic health. *Compend Contin Educ Dent* 2001;22:12-8.
 10. Faul F, Erdfelder E, Buchner A, Lang AG. Statistical power analyses using G* Power 3.1: Tests for correlation and regression analyses. *Behavior research methods*. 2009 Nov; 41(4):1149-60.
 11. Tonetti MS, Greenwell H, Kornman KS. Staging and grading of periodontitis: Framework and proposal of a new classification and case definition. *J Periodontol*. 2018;89(February):S159–72.
 12. Robert E. Schifferle. Periodontal disease and nutrition: separating the evidence from current fads. *Periodontology* 2000, Vol. 50, 2009, 78–89. doi:10.1111/j.1600-0757.2008.00297.x.
 13. Omare Mo, Kibet Jk, Cherutoi Jk, Kengara Fo. A Review Of Tobacco Abuse And Its Epidemiological Consequences. *Z Gesundheitswissenschaften*. 2022;30(6):1485–500.
 14. Rahul KL, Gauri S, James S, Miyola CF. Comparative Evaluation Of Serum Folic Acid, Vitamin B12 Levels And Erythrocyte Sedimentation Rate (Esr) In Smokers And Non-Smokers With Chronic Periodontitis: A Clinico-Biochemical Study, 2024; *Journal of Dental and Medical Sciences* e-ISSN: 2279-0853, p-ISSN: 2279-0861. 23, (1)PP 17-25.
 15. Alwan AM, Al Karawi SI, Abdul Wahab GA. The impact of traditional smokers versus electronic cigarette smokers on periodontal parameters in iraqi males. *Braz Dent Sci*. 2024;27(1):e4166. <https://doi.org/10.4322/bds.2024.e4166> .
 16. Singh A, Ladusingh L. Prevalence And Determinants Of Tobacco Use In India: Evidence From Recent Global Adult Tobacco Survey Data. *Plos One*. 2014;9(12):E114073.
 17. Torrungruang K, Tamsailom S, Rojanasomsith K, Sutdhibhisal S, Nisapakultorn K, Vanichjakvong O, Et Al. Risk Indicators Of Periodontal Disease In Older Thai Adults. *J Periodontol*. 2005 Apr;76(4):558–65.
 18. Haffajee Ad, Socransky Ss. Relationship Of Cigarette Smoking To Attachment Level Profiles. *J Clin Periodontol*. 2001 Apr;28(4):283–95.
 19. Calsina G, Ramón Jm, Echeverría Jj. Effects Of Smoking On Periodontal Tissues. *J Clin Periodontol*. 2002 Aug;29(8):771–6.
 20. Jalayer Naderi N, Semyari H, Elahinia Z. The Impact Of Smoking On Gingiva: A Histopathological Study. *Iran J Pathol*. 2015 Jul 1;10(3):214–20.
 21. Johannsen A, Asberg M, Söder Po, Söder B. Anxiety, Gingival Inflammation And Periodontal Disease In Non-Smokers And Smokers - An Epidemiological Study. *J Clin Periodontol*. 2005 May;32(5):488–91.
 22. Arowojolu Mo, Fawole Oi, Dosumu Eb, Opeodu Oi. A Comparative Study Of The Oral Hygiene Status Of Smokers And Non- Smokers In Ibadan, Oyo State. *Niger Med J J Niger Med Assoc*. 2013;54(4):240–3.

23. Ragghianti Ms, Greggi Sla, Lauris Jrp, Sant'ana Acp, Passanezi E. Influence Of Age, Sex, Plaque And Smoking On Periodontal Conditions In A Population From Bauru, Brazil. *J Appl Oral Sci Rev Fob*. 2004 Dec;12(4):273–9.
24. Velidandla S, Bodduru R, Birra V, Jain Y, Valluri R, Ealla Kkr. Distribution Of Periodontal Pockets Among Smokers And Nonsmokers In Patients With Chronic Periodontitis: A Cross-Sectional Study. *Cureus*. 2019 Sep 6;11(9):E5586.
25. Susin C, Haas An, Oppermann Rv, Haugejorden O, Albandar Jm. Gingival Recession: Epidemiology And Risk Indicators In A Representative Urban Brazilian Population. *J Periodontol*. 2004 Oct;75(10):1377–86.
26. Rudziński R, Banach J. [The Effect Of Tobacco Smoke On The Course And Severity Of Inflammation In Periodontal Tissues]. *Ann Acad Med Stetin*. 2011;57(1):88–95.
27. M Esaki, M Morita, R Akhter, K Akino, O Honda. Relationship between folic acid intake and gingival health in non-smoking adults in Japan. *Oral Diseases* Volume 16, Issue 1, pages 96–101, January 2010 doi:10.1111/j.1601-0825.2009.01619.x. PMID:19732352.
28. Vogel R and M Deasy. Folic acid and experimentally produced gingivitis. *IADR abstracts* 1977.
29. Ainamo J and Bay I 1975. Problems and proposals for recording gingivitis and plaque. *International Dental Journal* 25, 229-235. PMID:1058834.
30. Newman MG, Takei HH and Carranza FA. *Clinical Periodontology* 10th edition; Saunders.
31. Robert E. Schifferle. Periodontal disease and nutrition: separating the evidence from current fads. *Periodontology* 2000, V ol. 50, 2009, 78–89. doi:10.1111/j.1600-0757.2008.00297.x. PMID:19388955.
32. Neiva RF, Steigenga J, Al-Shammari KF, Wang H-L. Effects of specific nutrients on periodontal disease onset, progression and treatment. *J Clin Periodontol* 2003; 30: 579–589. doi:10.1034/j.1600-051X.2003.00354.x. PMID:12834494.
33. Erdemir EO, Bergstrom J. Effect of smoking on folic acid and vitamin B12 after nonsurgical periodontal intervention. *J Clin Periodontol* 2007; 34: 1074–1081. doi:10.1111/j.1600-051X.2007.01154.x. PMID:17953692.
34. Sumona, B., Sheetal, S., Anil, M., & Suvarna, P. (2011). Comparative evaluation of serum folic acid levels in smokers and non-smokers with chronic periodontitis. *Bangladesh Journal of Medical Science*, 10(2), 83–90.
35. Yu Yh, Kuo Hk, Lai Yl. The Association Between Serum Folate Levels And Periodontal Disease In Older Adults: Data From The National Health And Nutrition Examination Survey 2001/02. *J Am Geriatr Soc*. 2007 Jan;55(1):108–13.