

Scardovia wiggisiae as Recent Player in Caries Process: A Review

Raafat Jasim Al-Adool¹, Suhad Muwafaq Hamdoon¹ and Heba Faiz Hamodat¹

¹ Department of Prosthetic Dental Techniques, College of Health and Medical Techniques, Middle Technical University, Baghdad, Iraq

Correspondence: Raafat Jasim Al-Adool

Email: rafitadool2017@uomosul.edu.iq

Received: 21 April 2023; Accepted: 20 June 2023; Published: 28 June 2023

Abstract

Aim of the study: Oral cavity is attacked by many infectious agents that invade tooth, periodontium and tongue. Dental caries which is the most common oral disease, can be caused by the reaction between tooth structure, oral bacteria and fermentable carbohydrates. Most of caries microbial researches considered *Streptococcus mutans* and lactobacilli as major caries-associated players. The discovery of *Scardovia wiggisiae* in 2011 highlighted the possible occurrence of dental decay in absence of both *Streptococcus mutans* and lactobacilli in addition to other oral diseases. This review focused on taxonomy, acidogenicity, normal habitat and the important roles of newly *S.wiggisiae* as the pioneer bacteria for initiation and progression of caries process.

Material and method: The database was focused on articles published till 2022 for the topics oral diseases, dental caries, *Scardovia wiggisiae* and microbiology of dental caries.

Conclusion: *Scardovia wiggisiae* can be considered as a potential risk indicator for caries prediction in a given population.

Keywords: *Scardovia wiggisiae*, oral diseases, dental caries, microbiology of dental caries.

Introduction

Human kind faced oral infections since the human existence⁽¹⁾. Bacteria can play an important role in the initiation and later development of oral diseases⁽²⁾. Dental caries which is a common global problem⁽³⁾, has a complex infectious process with different causes⁽⁴⁾. It has major three risk factors; cariogenic bacteria, susceptible teeth, and the frequent dietary carbohydrates intake⁽⁵⁾. As the mouth has one of the most larger and most diverse microbiology⁽⁶⁾, understanding of diverse microbiota of dental caries is necessary because it is not a simple process caused by single bacteria but by a complicated tooth-bacterial interaction⁽⁷⁾.

Many researches have claimed that *Streptococcus mutans* is the primary major pathogen to initiate caries⁽⁸⁾ while others found these bacteria on dental surfaces; however, with no participating role in the development of caries lesion⁽⁹⁾.

Although *Streptococcus mutans* and lactobacilli are identified as caries-associated bacteria. It was noticed that other bacteria can cause dental caries in absence of the above bacteria, opening the door for discovering a new cariogenic bacteria⁽¹⁰⁾. The aim of this review was to identify the role of *Scardovia wiggisiae* in dental caries.

Materials and Methods

The Medline database was focused for English-language articles, published up to 2022 for the topics oral diseases, dental caries, *Scardovia wiggisiae* and microbiology of dental caries. Manual searching was additionally done from the citations of the updated identified reports. The literature search intent was done to highlight and review the analytic reports of the emerging and pioneer role of *Scardovia wiggisiae* in



many oral diseases and aspects include all peer-reviewed epidemiological and supporting human scientific articles.

Exclusion criteria: Studies that were reported in languages other than English were excluded due to practical difficulty of translation. In addition, articles that dealt with systemic effects, presence and diseases of *Scardovia wiggisiae* other than oral cavity were excluded.

Results and discussion

Tanner *et al* in 2011 applied updating molecular techniques to identified a new causative agent of dental caries in absence of known previous agents (e.g. *S. mutans*). This new agent was known as *Scardovia wiggisiae*. It should be noted that *Scardovia* was derived from the name of Italian microbiologist Vittorio Scardovi while *wiggisiae* was derived from American microbiologist Lois Wiggs^(11,12).

Scardovia belongs to Bifidobacteriaceae family that have seven genera, and due to presence of great alterations in genome sequence, *Scardovia* became out of the genus *Bifidobacterium*⁽¹³⁾. The systemic classification of *Scardovia wiggisiae* is

“Domain: Bacteria

Phylum: Actinobacteria

Class: Actinobacteridae

Order: Bifidobacteriales

Family: Bifidobacteriaceae

Genus: *Scardovia*

Species: *wiggisiae*^(14,15)”

Morphologically, *S.wiggisiae* is Gram positive anaerobic bacillus, non-spore forming, non-motile with dimensions of about 0.65 µm width and 1.6 to 4 µm length. It's shape can be pleomorphic, tend to curved slightly with club-shaped arrangement. It can be found singularly, in pairs or in short chains⁽¹⁶⁾.

In biochemical caries formation, one of the most important factors to initiate caries is capacity of cariogenic bacteria to build up a biofilm⁽¹⁷⁾ nevertheless, not all the oral bacteria can take part with the same pattern in the caries process⁽¹⁸⁾. The high metabolic rate, self protection from saliva and oral environments and ability to form organic acid that demineralize the hard tooth structure are important factors for cariogenic bacteria to grow within a biofilm⁽¹⁹⁾.

Regarding to these points, *S.wiggisiae* was presented as important caries associated bacteria due to it's ability to build up and acidification of the oral plaque^(20,21). It is hetero-fermentative bacteria that can use different carbohydrates forming lactate and acetic acids as fermentation final end products⁽²²⁾.

Accordingly, *S.wiggisiae* can be considered as high acidogenic and aciduric bacteria as *S.mutans*⁽²³⁾, however with different metabolic pathway (called fructose-6-phosphate) from that of *S.mutans* glycolytic metabolic pathway⁽²⁴⁾. This enables *S.wiggisiae* to be to more acid tolerant

and a high ecological competitive in low pH environments⁽²³⁾. Furthermore, it has a saccharolytic hydrolyzing ability by producing a lot amounts of acetic acids and little-moderate of lactic and formic acids from breaking sugar molecules e.g.sucrose, glucose, fructose and so on^(16,23). These large amounts of acetic acids have great chance to invade and decalcify enamel more than lactic acids⁽²⁵⁾. As result of it's advance acidurcity and acidogency which can decrease the environmental pH up to 3.5⁽²³⁾, *S.wiggisiae* can be considered as important pathogen in caries process together with other cariogenic players⁽¹⁰⁾.

The normal habitat of the Bifidobacteriaceae family to which *S. wiggisiae* belongs, is present primarily in human and animals gastro-intestinal tracts⁽²⁶⁾. Orally, *S.wiggisiae* is present in the saliva of children with early childhood caries and extensive early childhood caries⁽¹⁰⁾. A study done by Streiff *et al* 2017, insured the presence of *S.wiggisiae* DNA in the saliva of about 20 %of adults and 25% of pediatric patients⁽²⁶⁾. Regarding counting *S.wiggisiae* in oral cavity, Zhou J *et al* 2016 found it in less than 0.1% of the oral microbiota; however, it can seriously influenced oral community leading to dysbiosis⁽²⁷⁾. Nevertheless, Isaac R D *et al* 2022, noticed that *S.wiggisiae* can be one of the predominant bacteria in caries process comparing to *mutans streptococci* and *lactobacilli*⁽²⁸⁾. Periodontaly, *S.wiggisiae* can be found in gingival cervices, gingival crevicular fluids⁽²⁹⁾, supragingival plaque sites as in the buccal region to upper molar area and lingually to lower incisor ,but to

lower extent on ventral surface to the tongue⁽³⁰⁾.

Pathologically, Streiff BJ *et al* 2015 believed that the cause of moderate childhood gingivitis and adults periodontal pockets was due to the presence of *S.wiggisiae*⁽²⁶⁾. Nevertheless, Chakrapani *et al* 2019, showed a lower concentration of *S.wiggisiae* in subgingival biofilm in both healthy patients and in gingivitis and chronic periodontitis patients⁽³¹⁾.

In dental caries, *S.wiggisiae* usually occurs in initial enamel caries⁽³²⁾ affecting about 30% of children in the United States of America⁽¹¹⁾. In addition, it was found in active, deep and soft dentinal lesions of both dentitions, deciduous and permanent teeth,⁽¹⁵⁾and greatly in infected pulpal tissue of children⁽³³⁾, explaining the ability of this bacteria to be active player in dental caries alone. *S.wiggisiae* also was present in root carious lesions and can reach up to 6% of it's total microbiom⁽³⁴⁾. Nevertheless *S.wiggisiae* can be more in dentinal lesions than plaque samples⁽³⁵⁾. The presence of extra carious risks factors as fixed orthodontics appliances, may increase the prevalence of *S.wiggisiae* as they increase plaque accumulation and interfere with normal salivary self-cleaning⁽³⁶⁾.

S.wiggisiae can be found in higher rate in both adult⁽²⁶⁾ and pediatric patients using fixed orthodontic appliances⁽³⁷⁾. According to Mc Daniel *et al* 2021, more than 44 % pediatric orthodontic patients have *S.wiggisiae* in their samples⁽²⁹⁾.

In relation to smoking, *S.wiggisiae* has the ability to form more biofilms necessary for

caries lesions in smoker compared to non-smokers patients⁽³⁸⁾. This can be explained as nicotine can exaggerate the cariogenicity of *S.wiggisiae* .⁽³⁹⁾

Isolation of *S.wiggisiae* needs an anaerobic culture conditions⁽⁴⁰⁾ with pH 7, 5.5 till 5 that can obtains on neutral to acidic environments⁽¹¹⁾. Downes J. *et al* (2011) modified a new selective media with acid indicator to isolate *S.wiggisiae* effectively called *S.wiggisiae* selective medium(SWSM)⁽¹⁶⁾. Nevertheless, polymerase chain reaction (PCR) can be the cornerstone for isolation of *S.wiggisiae* using the known primers⁽⁴¹⁾.

Conclusions: There was an evidence that *S.wiggisiae* could be considered an efficient cariogenic, acidogenic and acidruic oral bacteria that has many criteria, enabling to be used as a potential risk indicator for caries prediction in a given population. This would provide a useful information for future development of new strategies of caries diagnosis, prevention and treatment.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Funding

This research received no external funding.

Data Availability Statement

Data are available from the authors upon reasonable request.

Acknowledgments

The authors would like to thank College of Dentistry, Mosul University, Mosul, Iraq, for their support in the present work.

References

1. Huang TT, Tseng FY, Yeh TH, Hsu CJ, Chen YS. Factors affecting the bacteriology of deep neck infection: A retrospective study of 128 patients. *Acta Otolaryngol.* 2006;126(4):396-401. <https://doi.org/10.1080/00016480500395195>
2. Bascones Martínez A, Figuero Ruiz E. Periodontal diseases as bacterial infection. *Av Periodon Implantol.* 2005;17(3):111-118. <https://doi.org/10.4321/S1699-65852005000300002>
3. Niederman R, Huang SS, Trescher AL, Listl S. Improving Oral Health Equity With Universal School-Based Caries Prevention. *Am J Public Health.* 2017;107(1):50-55. <https://doi.org/10.2105/AJPH.2016.303614>
4. Berkowitz RJ. Causes, treatment and prevention of early childhood caries: A microbiologic perspective. *J Can Dent Assoc.* 2003;69(5):304-307.
5. Tanner ACR. Anaerobic culture to detect periodontal and caries pathogens. *J Oral Biosci.* 2015;57(1):18-26. <https://doi.org/10.1016/j.job.2014.08.001>
6. Deo PN, Deshmukh R. Oral microbiome: Unveiling the fundamentals. *J Oral Maxillofac Pathol.* 2019;23(1):122-128. https://doi.org/10.4103/jomfp.JOMFP_304_18

7. Aas JA, Griffen AL, Dardis SR, Lee AM, Olsen I, Dewhirst FE, Leys EJ, Paster BJ. Bacteria of dental caries in primary and permanent teeth in children and young adults. *J Clin Microbiol.* 2008;46(4):1407-1417. <https://doi.org/10.1128/JCM.01410-07>
8. Loesche WJ, Rowan J, Straffon LH, Loos PJ. Association of Streptococcus mutans with human dental decay. *Infect Immun.* 1975;11(6):1252-1260. <https://doi.org/10.1128/iai.11.6.1252-1260.1975>
9. Sansone C, van Houte J, Joshipura K, Kent R, Margolis HC. The association of mutans streptococci and non-mutans streptococci capable of acidogenesis at a low pH with dental caries on enamel and root surfaces. *J Dent Res.* 1993;72(2):508-516. <https://doi.org/10.1177/00220345930720020701>
10. Chandna P, Srivastava N, Sharma A, Sharma V, Gupta N, Adlakha VK. Isolation of Scardovia wiggsiae using real-time polymerase chain reaction from the saliva of children with early childhood caries. *J Indian Soc Pedod Prev Dent.* 2018;36(3):290-295. https://doi.org/10.4103/JISPPD.JISPPD_225_17
11. Tanner AC, Mathney JM, Kent RL, Chalmers NI, Hughes CV, Loo CY, Pradhan N, Kanasi E, Hwang J, Dahlan M, Papadopoulou E. Cultivable anaerobic microbiota of severe early childhood caries. *J Clin Microbiol.* 2011;49(4):1464-1474. <https://doi.org/10.1128/JCM.02427-10>
12. Tanner AC, Kent RL, Holgerson PL, Hughes CV, Loo CY, Kanasi E, Chalmers NI, Johansson I. Microbiota of severe early childhood caries before and after therapy. *J Dent Res.* 2011;90(11):1298-1305. <https://doi.org/10.1177/0022034511421201>
13. Jian W, Dong X. Transfer of Bifidobacterium inopinatum and Bifidobacterium denticolens to Scardovia inopinata gen. nov., comb. nov., and Parascardovia denticolens gen. nov., comb. nov., respectively. *Int J Syst Evol Microbiol.* 2002;52(Pt 3):809-812. <https://doi.org/10.5802/aif.1902>
14. Mantzourani M, Gilbert SC, Sulong HN, Sheehy EC, Tank S, Fenlon M, Beighton D. The isolation of bifidobacteria from occlusal carious lesions in children and adults. *Caries Res.* 2009;43(4):308-313. <https://doi.org/10.1159/000222659>
15. Mantzourani M, Fenlon M, Beighton D. Association between Bifidobacteriaceae and the clinical severity of root caries lesions. *Oral Microbiol Immunol.* 2009;24(1):32-37. <https://doi.org/10.1111/j.1399-302X.2008.00470.x>
16. Downes J, Mantzourani M, Beighton D, Hooper S, Wilson MJ, Nicholson A, Wade WG. Scardovia wiggsiae sp. nov., isolated from the human oral cavity and clinical material, and emended descriptions of the genus Scardovia and Scardovia inopinata. *Int J Syst Evol Microbiol.* 2011;61(Pt 1):25-29. <https://doi.org/10.1099/ijs.0.019752-0>
17. Larsen T, Fiehn NE. Dental biofilm infections - an update. *APMIS.* 2017;125(4):376-384. <https://doi.org/10.1111/apm.12688>
18. Jenkinson HF. Beyond the oral microbiome. *Environ Microbiol.* 2011;13(12):3077-3087. <https://doi.org/10.1111/j.1462-2920.2011.02573.x>
19. Seow WK. Biological mechanisms of early childhood caries. *Community Dent Oral*

- Epidemiol. 1998;26(1 Suppl):8-27. <https://doi.org/10.1111/j.1600-0528.1998.tb02090.x>
20. Henne K, Rheinberg A, Conrads G, Esteves-Oliveira M. Analysis of Bacterial Activity in Sound and Cariogenic Biofilm: A Pilot in vivo Study. *Caries Res.* 2016;50(5):480-488. <https://doi.org/10.1159/000448485>
21. Hajishengallis E, Parsaei Y, Klein MI, Koo H. Advances in the microbial etiology and pathogenesis of early childhood caries. *Mol Oral Microbiol.* 2017;32(1):24-34. <https://doi.org/10.1111/omi.12152>
22. Pokusaeva K, Fitzgerald GF, van Sinderen D. Carbohydrate metabolism in Bifidobacteria. *Genes Nutr.* 2011;6(3):285-306. <https://doi.org/10.1007/s12263-010-0206-6>
23. Kameda M, Abiko Y, Washio J, Tanner ACR, Kressirer CA, Mizoguchi I, Takahashi N. Sugar Metabolism of *Scardovia wiggisiae*, a Novel Caries-Associated Bacterium. *Front Microbiol.* 2020;11:479. <https://doi.org/10.3389/fmicb.2020.00479>
24. Manome A, Abiko Y, Kawashima J, Washio J, Fukumoto S, Takahashi N. Acidogenic potential of oral Bifidobacterium and its high fluoride tolerance. *Front Microbiol.* 2019;10:1099. <https://doi.org/10.3389/fmicb.2019.01099>
25. Hoppenbrouwers PM, Driessens FC. The effect of lactic and acetic acid on the formation of artificial caries lesions. *J Dent Res.* 1988;67(12):1466-1467. <https://doi.org/10.1177/00220345880670120501>
26. Streiff BJ, Seneviratne M, Kingsley K. Screening and prevalence of the novel cariogenic pathogen *Scardovia wiggisiae* among adult orthodontic and non-orthodontic patient saliva samples. *Int J Dent Oral Health.* 2015;1(6). <https://doi.org/10.16966/2378-7090.159>
27. Zhou J, Jiang N, Wang S, Hu X, Jiao K, He X, et al. Exploration of human salivary microbiomes: insights into the novel characteristics of microbial community structure in caries and caries-free subjects. *PLoS One.* 2016;11(1):e0147039. <https://doi.org/10.1371/journal.pone.0147039>
28. Isaac RD, Sanjeev K, Subbulakshmi CL, Amirtharaj LV, Sekar M. Identification of a novel bacterium *Scardovia wiggisiae* in high caries risk adolescence: A metagenomic and melt curve analysis. *J Conserv Dent.* 2022;25(3):297-305.
29. McDaniel J, Howard KM, Kingsley K. Molecular Screening and Analysis Reveal Novel Oral Site-Specific Locations for the Cariogenic Pathogen *Scardovia wiggisiae*. *Dent J (Basel).* 2021;9(6):73. <https://doi.org/10.3390/dj906073>
30. Chervinets VM, Chervinets YV, Leont'Eva AV, Kozlova EA, Stulov NM, Belyaev VS, Grigoryants EO, Mironov AY. The microbiome of oral cavity patients with periodontitis, adhesive and biofilm forming properties. *Klin Lab Diagn.* 2021;66(1):45-51. <https://doi.org/10.18821/0869-2084-2021-66-1-45-51>
31. Chakrapani AR, Ajjappa BG, Prakash S, Bhat K. *Scardovia wiggisiae*: Role in Periodontal Health and Disease - A Pilot Study. *Indian J Dent Adv.* 2019;11(4):119-124.

32. Torlakovic L, Klepac-Ceraj V, Ogaard B, Cotton SL, Paster BJ, Olsen I. Microbial community succession on developing lesions on human enamel. *J Oral Microbiol.* 2012;4. <https://doi.org/10.3402/jom.v4i0.16125>
33. Ledezma-Rasillo G, Flores-Reyes H, Gonzalez-Amaro AM, Garrocho-Rangel A, Ruiz-Rodriguez MDS, Pozos-Guillen AJ. Identification of cultivable microorganisms from primary teeth with necrotic pulps. *J Clin Pediatr Dent.* 2010;34(4):329-333. <https://doi.org/10.17796/jcpd.34.4.20124lu111544377>
34. Damé-Teixeira N, Parolo CCF, Malz M, Devine DA, Do T. Gene expression profile of *Scardovia* spp. in the meta-transcriptome of root caries. *Braz Oral Res.* 2020;34:e0042. <https://doi.org/10.1590/1807-3107bor-2020.vol34.0042>
35. Vacharaksa A, Suvansopee P, Opaswanich N, Sukarawan W. PCR detection of *Scardovia wiggsiae* in combination with *Streptococcus mutans* for early childhood caries-risk prediction. *Eur J Oral Sci.* 2015;123(5):312-318. <https://doi.org/10.1111/eos.12208>
36. Maret D, Marchal-Sixou C, Vergnes JN, Hamel O, Georgelin-Gurgel M, Van Der Sluis L, Sixou M. Effect of fixed orthodontic appliances on salivary microbial parameters at 6 months: a controlled observational study. *J Appl Oral Sci.* 2014;22(1):38-43. <https://doi.org/10.1590/1678-775720130318>
37. El-Ezmerli NF, Gregory RL. Effect of nicotine on biofilm formation of *Streptococcus mutans* isolates from smoking and non-smoking subjects. *J Oral Microbiol.* 2019;11(1):1662275. <https://doi.org/10.1080/20002297.2019.1662275>
38. Balhaddad AA, Melo MAS, Gregory RL. Inhibition of nicotine-induced *Streptococcus mutans* biofilm formation by salts solutions intended for mouthrinses. *Restor Dent Endod.* 2019;44(2):e4. <https://doi.org/10.5395/rde.2019.44.e4>
39. Milne W, Rezaei G, Whiteley A, Kingsley K. Cariogenic Pathogen *Scardovia Wiggsiae* Screening Among Pediatric Orthodontic Patients: A Pilot Study. *Curr Res Dent.* 2018;9(1):1-5. <https://doi.org/10.3844/crdsp.2018.1.5>
40. Tanner AC, Sonis AL, Lif Holgerson P, Starr JR, Nunez Y, Kressirer CA, Paster BJ, Johansson I. White-spot lesions and gingivitis microbiotas in orthodontic patients. *J Dent Res.* 2012;91(9):853-858. <https://doi.org/10.1177/0022034512455031>